

Variability in imaging spectroscopy data across aspen stands is due to differences in genotype and foliar biochemistry.



Phil Townsend¹, Mike Madritch², Clayton Kingdon¹,
Aditya Singh¹, Karen Mock³ and Rick Lindroth¹

¹University of Wisconsin, ²Appalachian State University, ³Utah State University



Source: USGS



Source: Wikimedia Commons, photo by Mark Muir



Forest Ecology and Management

Available online 29 January 2013

In Press, Corrected Proof — Note to users



Recent declines of *Populus tremuloides* in North America linked to climate

James J. Worrall^a,  , Gerald E. Rehfeldt^b, Andreas Hamann^c, Edward H. Hogg^d, Suzanne B. Marchetti^a, Michael Michaelian^d, Laura K. Gray^c

^a US Forest Service, Rocky Mountain Region, Gunnison, CO 81230, USA

^b US Forest Service, Rocky Mountain Research Station, Moscow, ID 83843, USA

^c University of Alberta, Dept. of Renewable Resources, Edmonton, Alberta, Canada T6G 2H1

^d Canadian Forest Service, Northern Forestry Centre, Edmonton, Alberta, Canada T6H 3S5





Upper Midwest

Sampled and Imaged in 2009

- 33 Stands
- 391 unique genets
- 733 Trees
- 137 multi-ramet genets
- 19 Genets on multiple AVIRIS pixels (26)

Rocky Mountains

Sampled and Imaged in 2010

- 39 Stands
- 132 unique genets
- 941 Trees
- 57 multi-ramet genets
- 31 Genets on multiple AVIRIS pixels (41)



© 2012 Google

Google earth

154 m

25

F307
F306 F310
F309 F308
F323 F324 F302 F319
F325 F322 F321 F303 F317
F301 F305 F320
F315 F312 F318
F311 F313





F107 F110
F109 F106
F108
F121 F101 F102
F123 F122 F105 F117
F124 F125 F115 F120
F111 F103 F116
F113 F112 F114
F210 F208
F206 F207
F209
F221 F225
F202 F216 F217
F223 F201 F220
F214 F203 F212 F218
F215 F211

Foliar Nitrogen

-photosynthetic capacity

Highly variable in aspen:

1.13%-3.72%



Foliar Lignin

-regulates liquid transport
-recalcitrant (decomposition)

1.47%-22.21%



Condensed Tannins

-secondary metabolite



-variability not fully understood

0.20%-30.34%





- Strongly influence belowground processes
- Affect soil microbes (decrease respiration)
- Slow decomposition
- Reduce N mineralization rates
- Chelate metals




Variations in Foliar Chemistry




Mapping Foliar Chemistry with AVIRIS







Fish Lake National Forest


Fish Lake National Forest




August 2010



Red = Nitrogen Green = CT's



Red = Lignin Blue = Nitrogen





Ottawa National Forest


Northern
Wisconsin



July 2009





Red = Nitrogen Green = CT's



Red = Lignin Blue = Nitrogen

Spectral variability between genets:




Are genets distinguishable by spectra?


- Leaf traits correspond with traits related to belowground microbial function (and therefore spectra)
- Foliar and soil traits differ among genets, with foliar traits being more separable.
- Genets can be discriminated on the basis of leaf traits alone, but accuracies are much higher when using imaging spectroscopy.

Linkage to Soil Microbial Dynamics

- Biodiversity occurs at all levels of ecological organization (community, species, genetic, functional [chemical], etc.)
- Belowground processes:



Linkage to Belowground Function




Illumina bacterial sequencing
(pyrosequencing vs PLFA)
515f/806R primers

NMS
3 axis explain ~ 95% variation
in known bacterial classes

MRPP
T = -3.45
A = 0.16
P = 0.0016

- next gen sequencing of soil microbes, bacterial/fungal DNA
- construct barcode of barcoded primers
- match sequences against a known library of organisms
- hundreds of species that are unknown (larger w/ finer taxon)
- unknown kingdom is very very small (<0.01%),
- the percentage of unknown genus ID's is large (75%)
- we used class (35-40% unknown)

Summary of Project Findings



- Soil microbial function is related to canopy chemistry
- Genotypes have distinct soil microbial profiles
- AVIRIS imagery predicts canopy chemistry: lignin and CTs
- Genotypes are spectrally distinct
- AVIRIS imagery predicts variation in belowground soil function